

Dual R-Smads interplay in the regulation of vertebrate neurogenesis

Adriana L Sanz^{1,2,3} and David G Míguez^{1,2,3,*}

¹Depto. de Física de la Materia Condensada; Universidad Autónoma de Madrid; Campus de Cantoblanco; Madrid, Spain; ²Instituto de Materiales Nicolás Cabrera; Universidad Autónoma de Madrid; Campus de Cantoblanco; Madrid, Spain; ³Condensed Matter Physics Center IFIMAC; Universidad Autónoma de Madrid; Campus de Cantoblanco; Madrid, Spain

During neurogenesis, multiple regulatory networks integrate extracellular and intracellular signals to ensure proper final numbers for each neuronal subtype at the end of the developmental process. The Activin/TGF- β signaling cascade is one of the main players of neurogenesis in vertebrates, balancing proliferation and differentiation of neural stem cells by regulating gene expression via the R-Smads transcription factors. Despite their equivalent upstream activation mechanism, Smad2 and Smad3 functions can be redundant or opposite depending on the particular context of the cell. We demonstrate that R-Smad simultaneously cooperate and antagonize in the regulation of gene expression in the context of vertebrate neurogenesis. We propose a model where synergism and antagonism appear as a consequence of the competition between Smad2 and Smad3 to form the different transcriptionally active heterotrimers with Smad4. This dual interplay of R-Smads significantly modulates the role of TGF- β pathway in the balance between proliferation and differentiation in the developing vertebrate spinal cord.

arrest, differentiation and migration. The canonical TGF- β pathway begins with the binding of TGF- β ligands to different isoforms of type II and I Ser/Thr kinase receptors at the cell membrane.³ The ligand promotes receptor dimerization, enabling the phosphorylation and activation of the receptor type I by the type II. Specifically, activated Nodal/Activin/TGF- β receptor complexes recruit and phosphorylate Smad2 and Smad3 proteins, which oligomerize with Smad4 to form transcriptionally active heterotrimers. These active complexes translocate into the nucleus where they regulate gene expression (Fig. 1A).⁴⁻⁶

Despite this apparently simple mechanism of activation, the Nodal/Activin/TGF- β pathway plays major and diverse roles during embryo development. The concentration gradient of Nodal is essential for the establishment of the anterior-posterior (A/P) axis and the induction of the mesendoderm layer. Subsequently, together with BMPs, they organize the left-right (L/R) asymmetry, finishing the specification of the body plan.^{1,7} Counteracting BMP and Shh gradients regulate closure and dorsal-ventral (D/V) polarization of the neural tube.⁸⁻¹⁰ BMP/Activin ligands are expressed dorsally in the newly originated neural tube,¹¹ whereas TGF- β 2, 3 isoforms are preferentially expressed in the notochord and the floor plate,^{12,13} concomitant with Shh expression. The opposing gradients of these ligands are involved in the specification pattern for each neuronal subtype in the neural tube. Nonetheless, the particular functions of each member of the Activin/TGF- β subfamily

Keywords: neural tube, differentiation, networks, systems biology

*Correspondence to: David G Míguez;
Email: david.gomez.miguez@uam.es

Submitted: 03/07/2014

Revised: 05/21/2014

Accepted: 06/09/2014

Published Online: 06/12/2014

Citation: Sanz AL, Míguez DG. Dual R-Smads interplay in the regulation of vertebrate neurogenesis. *Neurogenesis* 2014; 1:e29529; <http://dx.doi.org/10.4161/neur.29529>

Commentary to: Míguez DG, Gil-Guiñón E, Pons S, Martí E. Smad2 and Smad3 cooperate and antagonize simultaneously in vertebrate neurogenesis. *J Cell Sci* 2013; 126:5335-43; PMID:24105267; <http://dx.doi.org/10.1242/jcs.130435>

during neurogenesis are beginning to be understood.

In the dorsal pole of the neural tube, Activin signaling promotes the formation of the dorsal interneuron 3 (dI3) precursor cells.¹⁴ Besides, TGF- β factors contribute to regulate the balance between proliferation and differentiation of neural stem cells in conjunction with other pathways. TGF- β counterbalances the proliferative signals of IGF1/AKT in the forebrain and Wnt/FGF8 in the midbrain.^{15,16} In addition, TGF- β 1 signaling participates in neuronal differentiation by modulating the expression of regulators of the cell cycle,^{17,18} whereas the isoforms TGF- β 2,3 are essential for the specification of dopaminergic neurons induced by Shh.¹⁹

The role of Activin/TGF- β signaling in the regulation of neurogenesis is ultimately mediated by the downstream transcription factors Smad2 and Smad3. Particularly, Smad3 triggers the differentiation program of ventral interneuron progenitors through transcription of bHLH proteins such as NeuroM and Mash1, inhibition of Id1,2,^{20,21} and promoting cell-cycle exit by transcribing cell-cycle inhibitors p15, p21, or p27.^{17,20,21} In contrast, Smad2 activity is correlated with Nodal signaling throughout embryogenesis, including A/P patterning and endoderm formation, although its specific function in vertebrate neurogenesis remains unclear.⁷ However, recent studies suggest a role of Smad2 in the regulation of axonal morphogenesis in telencephalon and metencephalon,^{22,23} which correlates with the pro-survival effects that TGF- β has in neurons in other contexts.^{24,25} The role of Smad2 in neurogenesis appears to be related to SnoN, a modulator of axonal development and growth.²⁶ The SnoN pathway is receiving increased importance during embryogenesis, since recent findings suggest a high convergence with other important pathways, including Nodal/TGF- β ²⁷ and cell cycle regulators such as p53 and APC/Cdh1.²⁸ Furthermore, in some cell lines, SnoN expression seems to be positively and negatively regulated by both R-Smads, revealing a potential implication of both R-Smads in this pathway.²⁹

Related to the expression profile, Smad2 is widely distributed in the neuroepithelium, and latterly in the telencephalon

and metencephalon.³⁰ Nonetheless, in the context of the developing neural tube, Smad2 was not detected in previous studies.²⁰ We showed that Smad2 is expressed in the ventricular zone of the developing spinal cord in the chick neural tube, and that its expression partially overlaps with the expression pattern of Smad3, which is a requirement for a potential interplay between both R-Smads at the cellular level.³¹

As reviewed above, Smad2 and Smad3 are functionally nonequivalent,^{5,32} despite their common upstream activation mechanism and their 92% identity in their amino acid sequences.³²⁻³⁴ Knockout (KO) studies in mice, show early embryonic lethality in Smad2 KO. This was attributed to disruption of the A/P axis within the epiblast and failure in the formation of the ectoderm, mesoderm, and endoderm.³⁴ In contrast, the KO of Smad3 is not embryonic lethal, although mice present deficiencies in the immune response and high predisposition to tumorigenesis.³⁵ In addition, Smad2 and Smad3 have been shown to recruit different co-factors and target different regulatory sequences.³⁶ In particular, Smad3 binds directly to a CAGA-box in the promoter of PAI-1 or c-Jun, whereas Smad2 binds indirectly through Smad4 and co-factors, such as FoxH1 and Mix families, to Activin response elements (ARE) or other regulatory sequences.^{33,37} Moreover, recent studies suggest that cell-type specific master transcription factors such as Oct4, Myod1, or PU.1 determine Smad2 and Smad3 targets, and hence direct TGF- β effects in a context-dependent manner.³⁸

Despite the important and multiple functions of the TGF- β signaling in many cellular processes and diseases,^{39,40} there are still relevant open questions regarding to the global mechanisms that govern TGF- β dynamics and function, with the 2 R-Smads playing redundant roles in some scenarios^{9,10} while antagonizing in others.^{41,42}

To answer the question of how these 2 highly similar molecules targeting different DNA regulatory sequences can synergize or antagonize depending on the cellular context, we compared the function of both R-Smads in the context of vertebrate neurogenesis.³¹ Our studies

revealed a simultaneous cooperation and antagonism between Smad2 and Smad3 occurring at the same cellular context. Experiments performed in the chick neural tube by *in ovo* electroporation showed an increase of Smad3-specific transcription targets after Smad2-Smad3 gain of function (GOF) compared with wild-type (WT) or Smad3 GOF alone. Surprisingly, Smad2 loss of function (LOF) also induced a strong increase of Smad3 activity (Fig. 1B). Additionally, Smad2-Smad3 GOF resulted in an increase of the amount of neural progenitors undergoing differentiation, specifically of ventral interneurons, compared with WT or GOF of Smad3.²⁰ Correspondingly, LOF of Smad2 also induced a strong increase in differentiation, compared with the WT situation.

In order to address the potential underlying mechanisms that could induce this dual interplay between the R-Smads, we followed a computational approach. Previous models of the TGF- β signaling pathway focused on the dynamics of the nucleocytoplasmic shuttling of R-Smad complexes,⁴³ the dynamics of signal processing, the generation of transient responses,⁴⁴ and endocytosis.⁴⁵ Most of these mathematical models assume that Smad2 and Smad3 are indistinguishable. These approaches take advantage of a detailed characterization of several potential interactions in the pathway to understand the effect of different regulatory mechanisms and crosstalks.⁴⁶

Contrary, our modeling approach focused on the R-Smads oligomerization after phosphorylation, in order to understand the dual antagonism/cooperation observed experimentally. This minimal model successfully reproduced all experimental results when we allowed the heterotrimer Smad2-Smad3-Smad4 to bind and regulate transcription in Smad3-specific regulatory sequences. We observed by co-immunoprecipitation that Smad2 and Smad3 physically interact in a complex with Smad4, and that the amount of Smad3 bound to Smad4 depended on the amount of Smad2 available and vice versa. The affinity of phosphorylated Smad2 to form trimers with Smad2 vs. Smad3 is unknown. Nonetheless, the model predicted stronger affinity of the R-Smads




Figure 1. Scheme of the TGF- β pathway. **(A)** (1) Dimers of soluble Activin or TGF- β ligands bind their specific Ser/Thr kinase receptor type II, presented by the co-receptor betaglycan. (2) The complex oligomerizes and phosphorylates receptor type I and finally forms the tetrameric active ligand-receptor complex. (3) This complex phosphorylates R-Smad transcription factors (Smad2 in green and Smad3 in blue), recruited to plasma membrane by the anchor protein SARA. (4) Heterodimers or homodimers of R-Smad bind the Co-Smad (Smad4) to form the transcriptionally active heterotrimers. (5) Smad oligomers are actively transported to the nucleus, where (6) they bind to specific promoter regions and regulate gene expression. Specifically, Smad2-Smad2-Smad4 homotrimer is recruited to ARE while Smad3-Smad3-Smad4 homotrimer and Smad3-Smad2-Smad4 heterotrimer bind to CAGA box in the promoter regions of their target genes. **(B)** In the basal state Smad2 directed transcription (ARE) is accomplished only by Smad2-Smad2-Smad4 homotrimer while Smad3 targets (CAGA box) are regulated by both Smad3-Smad2-Smad4 and Smad3-Smad3-Smad4 trimers. Increasing the levels of Smad2 (Smad2 GOF) shifts the balance to overexpress Smad2 targets (ARE). Similarly, Smad3 GOF leads to overexpression of Smad3 targets. Depletion of Smad2 protein (Smad2 LOF) contributes not only to downregulate Smad2 targets, but also to increase Smad3 driven transcription.

to form the heterotrimer Smad2-Smad3-Smad4, than Smad2-Smad2-Smad4 or Smad3-Smad3-Smad4.

Overall, our results evidence a scenario of competition between R-Smad to form the 3 potential heterotrimers that depends on the concentration and the binding affinities of the R-Smads. Thus, the selection between cooperation and antagonism of the R-Smads is not cell type-dependent, occurring in the same cellular context as a consequence of the interaction between Smad2, Smad3, and Smad4.

Many aspects of the TGF- β pathway were not included in the analysis for the sake of simplicity. In our model, we assumed Michaelis–Menten kinetics for the phosphorylation dynamics of the R-Smads, and therefore the amount of phosphorylated R-Smads depended directly on the amount of available R-Smads after receptor activation. Moreover, we considered that protein production and degradation are balanced. Consequently, the concentration of each R-Smad was assumed constant. A more detailed analysis of our model, including a full mathematical description of the known interactions of the TGF- β pathway, remains to be studied. Potential candidates to include in the model are compartmentalization, nucleo-cytoplasmic translocation of the R-Smads,⁴⁷ and inhibition by I-Smads (Smad6 and Smad7) and E3 ligases.⁴⁸ The I-Smads induce a well-characterized negative feedback loop that can induce nonlinear effects, such as desensitization of the pathway and robustness against perturbations.⁴⁹ Interestingly, some E3 ligases such as Arkadia have been demonstrated to positively contribute to TGF- β signaling. Arkadia has an essential role during embryogenesis because, under TGF- β activation, it degrades negative feedback inhibitors of the pathway, such as Smad7 and SnoN, a negative co-factor for Smad3/4 directed transcription.^{44,50} All these potential interactions will result in a more realistic mathematical approximation to the true dynamics of the pathway, but the increasing complexity and the amount of unknown reaction parameters will complicate the analysis and the understanding of the consequences of the interplay between the R-Smads.

The arising characteristics of the TGF- β system constitute a paradigm of how the network of interactions can determine the strength, dynamics and even the function of proteins in a given pathway. These emerging characteristics induced by a complex network have been studied widely in biological systems.⁵¹ In the context of neurogenesis, Shh graded responses have been proposed to be interpreted via a nonlinear network of interacting genes.⁸ In addition, domain specification of the neural tube has been also shown to be regulated by a network of interacting genes with multiple interconnected positive and negative feedback loops.⁵² For instance, a feedback in Delta-Notch signaling that mediates lateral inhibition is key in the regulation of pro-neural genes and neural differentiation.⁵³ In fact, the characterization of the interactions within the network has been shown to be decisive for a full understanding of different developmental processes like somite formation,⁵⁴ palate,⁵⁵ and digit formation in mice⁵⁶ and pigmentation pattern in animal skin⁵⁷ and also in treatment and prognosis of pathologies such as cancer.^{58–60} These studies evidence that a complete understanding of the TGF- β and other signaling networks involved in neurogenesis should take into account not only the study of the biochemical aspects of the proteins, but also the analysis of the interaction network.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

1. Wu MY, Hill CS. Tgf- β superfamily signaling in embryonic development and homeostasis. *Dev Cell* 2009; 16:329–43; PMID:19289080; <http://dx.doi.org/10.1016/j.devcel.2009.02.012>
2. Oshimori N, Fuchs E. The harmonies played by TGF- β in stem cell biology. *Cell Stem Cell* 2012; 11:751–64; PMID:23217421; <http://dx.doi.org/10.1016/j.stem.2012.11.001>
3. Feng XH, Derynck R. Specificity and versatility in tgf- β signaling through Smads. *Annu Rev Cell Dev Biol* 2005; 21:659–93; PMID:16212511; <http://dx.doi.org/10.1146/annurev.cellbio.21.022404.142018>
4. Heldin CHH, Miyazono K, ten Dijke P. TGF- β signalling from cell membrane to nucleus through SMAD proteins. *Nature* 1997; 390:465–71; PMID:9393997; <http://dx.doi.org/10.1038/37284>
5. Massagué J, Seoane J, Wotton D. Smad transcription factors. *Genes Dev* 2005; 19:2783–810; PMID:16322555; <http://dx.doi.org/10.1101/gad.1350705>
6. ten Dijke P, Hill CS. New insights into TGF- β -Smad signalling. *Trends Biochem Sci* 2004; 29:265–73; PMID:15130563; <http://dx.doi.org/10.1016/j.tibs.2004.03.008>
7. Brennan J, Lu CC, Norris DP, Rodriguez TA, Beddington RS, Robertson EJ. Nodal signalling in the epiblast patterns the early mouse embryo. *Nature* 2001; 411:965–9; PMID:11418863; <http://dx.doi.org/10.1038/35082103>
8. Dessaud E, McMahon AP, Briscoe J. Pattern formation in the vertebrate neural tube: a sonic hedgehog morphogen-regulated transcriptional network. *Development* 2008; 135:2489–503; PMID:18621990; <http://dx.doi.org/10.1242/dev.009324>
9. Jia S, Wu D, Xing C, Meng A. Smad2/3 activities are required for induction and patterning of the neuroectoderm in zebrafish. *Dev Biol* 2009; 333:273–84; PMID:19580801; <http://dx.doi.org/10.1016/j.ydbio.2009.06.037>
10. Jia S, Ren Z, Li X, Zheng Y, Meng A. smad2 and smad3 are required for mesendoderm induction by transforming growth factor-beta/nodal signals in zebrafish. *J Biol Chem* 2008; 283:2418–26; PMID:18025082; <http://dx.doi.org/10.1074/jbc.M707578200>
11. Liu A, Niswander LA. Bone morphogenetic protein signalling and vertebrate nervous system development. *Nat Rev Neurosci* 2005; 6:945–54; PMID:16340955; <http://dx.doi.org/10.1038/nrn1805>
12. Liem KFJ Jr., Tremml G, Jessell TM. A role for the roof plate and its resident TGFbeta-related proteins in neuronal patterning in the dorsal spinal cord. *Cell* 1997; 91:127–38; PMID:9335341; [http://dx.doi.org/10.1016/S0092-8674\(01\)80015-5](http://dx.doi.org/10.1016/S0092-8674(01)80015-5)
13. Cooley JR, Yatskevich TA, Antin PB. Embryonic expression of the transforming growth factor β ligand and receptor genes in chicken. *Dev Dyn* 2014; 243:497–508; PMID:24166734; <http://dx.doi.org/10.1002/dvdy.24085>
14. Timmer J, Chesnutt C, Niswander L. The activin signaling pathway promotes differentiation of dl3 interneurons in the spinal neural tube. *Dev Biol* 2005; 285:1–10; PMID:16039645; <http://dx.doi.org/10.1016/j.ydbio.2005.05.039>
15. Seoane J, Le HV, Shen L, Anderson SA, Massagué J. Integration of Smad and forkhead pathways in the control of neuroepithelial and glioblastoma cell proliferation. *Cell* 2004; 117:211–23; PMID:15084259; [http://dx.doi.org/10.1016/S0092-8674\(04\)00298-3](http://dx.doi.org/10.1016/S0092-8674(04)00298-3)
16. Falk S, Wurdak H, Ittner LM, Ille F, Sumara G, Schmid MT, Draganova K, Lang KS, Paratore C, Leveen P, et al. Brain area-specific effect of TGF- β signaling on Wnt-dependent neural stem cell expansion. *Cell Stem Cell* 2008; 2:472–83; PMID:18462697; <http://dx.doi.org/10.1016/j.stem.2008.03.006>
17. Misumi S, Kim TSS, Jung CG, Masuda T, Urakawa S, Isobe Y, Furuyama F, Nishino H, Hida H. Enhanced neurogenesis from neural progenitor cells with G1/S-phase cell cycle arrest is mediated by transforming growth factor β 1. *Eur J Neurosci* 2008; 28:1049–59; PMID:18783370; <http://dx.doi.org/10.1111/j.1460-9568.2008.06420.x>
18. Gomes FC, Sousa VdeO, Romão L. Emerging roles for TGF- β 1 in nervous system development. *Int J Dev Neurosci* 2005; 23:413–24; PMID:15936920; <http://dx.doi.org/10.1016/j.ijdevneu.2005.04.001>
19. Farkas LM, Dünker N, Roussa E, Unsicker K, Kriegstein K. Transforming growth factor- β (s) are essential for the development of midbrain dopaminergic neurons in vitro and in vivo. *J Neurosci* 2003; 23:5178–86; PMID:12832542

20. García-Campmany L, Martí E. The TGFbeta intracellular effector Smad3 regulates neuronal differentiation and cell fate specification in the developing spinal cord. *Development* 2007; 134:65-75; PMID:17138664; <http://dx.doi.org/10.1242/dev.02702>
21. Estarás C, Akizu N, García A, Beltrán S, de la Cruz X, Martínez-Balbás MA. Genome-wide analysis reveals that Smad3 and JMJD3 HDM co-activate the neural developmental program. *Development* 2012; 139:2681-91; PMID:22782721; <http://dx.doi.org/10.1242/dev.078345>
22. Ueberham U, Lange P, Ueberham E, Brückner MK, Hartlage-Rübsamen M, Pannicke T, Rohn S, Cross M, Arendt T. Smad2 isoforms are differentially expressed during mouse brain development and aging. *Int J Dev Neurosci* 2009; 27:501-10; PMID:19375497; <http://dx.doi.org/10.1016/j.ijdevneu.2009.04.001>
23. Wang L, Nomura M, Goto Y, Tanaka K, Sakamoto R, Abe I, Sakamoto S, Shibata A, Enciso PL, Adachi M, et al. Smad2 protein disruption in the central nervous system leads to aberrant cerebellar development and early postnatal ataxia in mice. *J Biol Chem* 2011; 286:18766-74; PMID:21464123; <http://dx.doi.org/10.1074/jbc.M111.223271>
24. Pineda JR, Daynac M, Chicheportiche A, Cebrian-Silla A, Sii Felice K, García-Verdugo JM, Boussin FD, Mouthon MA. Vascular-derived TGF-β increases in the stem cell niche and perturbs neurogenesis during aging and following irradiation in the adult mouse brain. *EMBO Mol Med* 2013; 5:548-62; PMID:23526803; <http://dx.doi.org/10.1002/emmm.201202197>
25. Brionne TC, Teseur I, Masliah E, Wyss-Coray T. Loss of TGF-β 1 leads to increased neuronal cell death and microgliosis in mouse brain. *Neuron* 2003; 40:1133-45; PMID:14687548; [http://dx.doi.org/10.1016/S0896-6273\(03\)00766-9](http://dx.doi.org/10.1016/S0896-6273(03)00766-9)
26. Stegmüller J, Huynh MA, Yuan Z, Konishi Y, Bonni A. TGFbeta-Smad2 signaling regulates the Cdh1-APC/SnoN pathway of axonal morphogenesis. *J Neurosci* 2008; 28:1961-9; PMID:18287512; <http://dx.doi.org/10.1523/JNEUROSCI.3061-07.2008>
27. Zhu Q, Kim YH, Wang D, Oh SP, Luo K. SnoN facilitates ALK1-Smad1/5 signaling during embryonic angiogenesis. *J Cell Biol* 2013; 202:937-50; PMID:24019535; <http://dx.doi.org/10.1083/jcb.201208113>
28. Zhu Q, Luo K. SnoN in regulation of embryonic development and tissue morphogenesis. *FEBS Lett* 2012; 586:1971-6; PMID:22710172; <http://dx.doi.org/10.1016/j.febslet.2012.03.005>
29. Zhu Q, Pearson-White S, Luo K. Requirement for the SnoN oncoprotein in transforming growth factor beta-induced oncogenic transformation of fibroblast cells. *Mol Cell Biol* 2005; 25:10731-44; PMID:16314499; <http://dx.doi.org/10.1128/MCB.25.24.10731-10744.2005>
30. de Sousa Lopes SM, Carvalho RL, van den Driesche S, Goumans MJ, ten Dijke P, Mummery CL. Distribution of phosphorylated Smad2 identifies target tissues of TGF β ligands in mouse development. *Gene Expr Patterns* 2003; 3:355-60; PMID:12799085; [http://dx.doi.org/10.1016/S1567-133X\(03\)00029-2](http://dx.doi.org/10.1016/S1567-133X(03)00029-2)
31. Míguez DG, Gil-Guiñón E, Pons S, Martí E. Smad2 and Smad3 cooperate and antagonize simultaneously in vertebrate neurogenesis. *J Cell Sci* 2013; 126:5335-43; PMID:24105267; <http://dx.doi.org/10.1242/jcs.130435>
32. Brown KA, Pietenpol JA, Moses HL. A tale of two proteins: differential roles and regulation of Smad2 and Smad3 in TGF-beta signaling. *J Cell Biochem* 2007; 101:9-33; PMID:17340614; <http://dx.doi.org/10.1002/jcb.21255>
33. Dennler S, Huet S, Gauthier JM. A short amino-acid sequence in MH1 domain is responsible for functional differences between Smad2 and Smad3. *Oncogene* 1999; 18:1643-8; PMID:10102636; <http://dx.doi.org/10.1038/sj.onc.1202729>
34. Dunn NR, Koone CH, Anderson DC, Islam A, Bikoff EK, Robertson EJ. Mice exclusively expressing the short isoform of Smad2 develop normally and are viable and fertile. *Genes Dev* 2005; 19:152-63; PMID:15630024; <http://dx.doi.org/10.1101/gad.1243205>
35. Ashcroft GS, Yang X, Glick AB, Weinstein M, Letterio JL, Mizel DE, Anzano M, Greenwell-Wild T, Wahl SM, Deng C, et al. Mice lacking Smad3 show accelerated wound healing and an impaired local inflammatory response. *Nat Cell Biol* 1999; 1:260-6; PMID:10559937; <http://dx.doi.org/10.1038/12971>
36. Morikawa M, Koinuma D, Miyazono K, Heldin CH. Genome-wide mechanisms of Smad binding. *Oncogene* 2013; 32:1609-15; PMID:22614010; <http://dx.doi.org/10.1038/ncr.2012.191>
37. Derynck R, Zhang Y, Feng XH. Smads: transcriptional activators of TGF-β responses. *Cell* 1998; 95:737-40; PMID:9865691; [http://dx.doi.org/10.1016/S0092-8674\(00\)81696-7](http://dx.doi.org/10.1016/S0092-8674(00)81696-7)
38. Mullen AC, Orlando DA, Newman JJ, Lovén J, Kumar RM, Bilodeau S, Reddy J, Guenther MG, DeKoter RP, Young RA. Master transcription factors determine cell-type-specific responses to TGF-β signaling. *Cell* 2011; 147:565-76; PMID:22036565; <http://dx.doi.org/10.1016/j.cell.2011.08.050>
39. Akhurst RJ, Hata A. Targeting the TGFβ signalling pathway in disease. *Nat Rev Drug Discov* 2012; 11:790-811; PMID:23000686; <http://dx.doi.org/10.1038/nrd3810>
40. Massagué J. TGFbeta in Cancer. *Cell* 2008; 134:215-30; PMID:18662538; <http://dx.doi.org/10.1016/j.cell.2008.07.001>
41. Labbé E, Silvestri C, Hoodless PA, Wrana JL, Attisano L. Smad2 and Smad3 positively and negatively regulate TGF β-dependent transcription through the forkhead DNA-binding protein FAST2. *Mol Cell* 1998; 2:109-20; PMID:9702197; [http://dx.doi.org/10.1016/S1097-2765\(00\)80119-7](http://dx.doi.org/10.1016/S1097-2765(00)80119-7)
42. Yang YCC, Piek E, Zavadil J, Liang D, Xie D, Heyer J, Pavlidis P, Kuchelapati R, Roberts AB, Böttinger EP. Hierarchical model of gene regulation by transforming growth factor β. *Proc Natl Acad Sci U S A* 2003; 100:10269-74; PMID:12930890; <http://dx.doi.org/10.1073/pnas.1834070100>
43. Schmierer B, Tournier AL, Bates PA, Hill CS. Mathematical modeling identifies Smad nucleocytoplasmic shuttling as a dynamic signal-interpreting system. *Proc Natl Acad Sci U S A* 2008; 105:6608-13; PMID:18443295; <http://dx.doi.org/10.1073/pnas.0710134105>
44. Nicklas D, Saiz L. Characterization of negative feedback network motifs in the TGF-β signaling pathway. *PLoS One* 2013; 8:e83531; PMID:24386222; <http://dx.doi.org/10.1371/journal.pone.0083531>
45. Cellière G, Fengos G, Hervé M, Iber D. Plasticity of TGF-β signaling. *BMC Syst Biol* 2011; 5:184; PMID:22051045; <http://dx.doi.org/10.1186/1752-0509-5-184>
46. Zi Z, Chapnick DA, Liu X. Dynamics of TGF-β/Smad signaling. *FEBS Lett* 2012; 586:1921-8; PMID:22710166; <http://dx.doi.org/10.1016/j.febslet.2012.03.063>
47. Inman GJ, Nicolás FJ, Hill CS. Nucleocytoplasmic shuttling of Smads 2, 3, and 4 permits sensing of TGF-beta receptor activity. *Mol Cell* 2002; 10:283-94; PMID:12191474; [http://dx.doi.org/10.1016/S1097-2765\(02\)00585-3](http://dx.doi.org/10.1016/S1097-2765(02)00585-3)
48. Saritas-Yildirim B, Silva EM. The role of targeted protein degradation in early neural development. *Genesis* 2014; 52:287-99; PMID:24623518; <http://dx.doi.org/10.1002/dvg.22771>
49. Míguez DG. Network nonlinearities in drug treatment. *Interdiscip Sci* 2013; 5:85-94; PMID:23740389; <http://dx.doi.org/10.1007/s12539-013-0165-x>
50. Levy L, Howell M, Das D, Harkin S, Episkopou V, Hill CS. Arkadia activates Smad3/Smad4-dependent transcription by triggering signal-induced SnoN degradation. *Mol Cell Biol* 2007; 27:6068-83; PMID:17591695; <http://dx.doi.org/10.1128/MCB.00664-07>
51. Oltvai ZN, Barabási AL. Systems biology. Life's complexity pyramid. *Science* 2002; 298:763-4; PMID:12399572; <http://dx.doi.org/10.1126/science.1078563>
52. Kicheva A, Cohen M, Briscoe J. Developmental pattern formation: insights from physics and biology. *Science* 2012; 338:210-2; PMID:23066071; <http://dx.doi.org/10.1126/science.1225182>
53. Formosa-Jordan P, Ibañes M, Ares S, Frade JM. Regulation of neuronal differentiation at the neurogenic wavefront. *Development* 2012; 139:2321-9; PMID:22669822; <http://dx.doi.org/10.1242/dev.076406>
54. Kaern M, Míguez DG, Muñozuri AP, Menzinger M. Control of chemical pattern formation by a clock-and-wavefront type mechanism. *Biophys Chem* 2004; 110:231-8; PMID:15228959; <http://dx.doi.org/10.1016/j.bpc.2004.02.006>
55. Economou AD, Ohazama A, Porntaveetus T, Sharpe PT, Kondo S, Basson MA, Gritli-Linde A, Coburn MT, Green JB. Periodic stripe formation by a Turing mechanism operating at growth zones in the mammalian palate. *Nat Genet* 2012; 44:348-51; PMID:22344222; <http://dx.doi.org/10.1038/ng.1090>
56. Sheth R, Marcon L, Bastida MF, Junco M, Quintana L, Dahn R, Kmita M, Sharpe J, Ros MA. Hox genes regulate digit patterning by controlling the wavelength of a Turing-type mechanism. *Science* 2012; 338:1476-80; PMID:23239739; <http://dx.doi.org/10.1126/science.1226804>
57. Míguez DG, Muñozuri AP. On the orientation of stripes in fish skin patterning. *Biophys Chem* 2006; 124:161-7; PMID:16844282; <http://dx.doi.org/10.1016/j.bpc.2006.06.014>
58. Míguez DG. The role of asymmetric binding in ligand-receptor systems with 1:2 interaction ratio. *Biophys Chem* 2010; 148:74-81; PMID:20332059; <http://dx.doi.org/10.1016/j.bpc.2010.02.012>
59. Doldán-Martelli V, Guantes R, Míguez DG. A mathematical model for the rational design of chimeric ligands in selective drug therapies. *CPT Pharmacometrics Syst Pharmacol* 2013; 2:e26; PMID:23887616; <http://dx.doi.org/10.1038/psp.2013.2>
60. Ruiz-Herrero T, Estrada J, Guantes R, Míguez DG. A tunable coarse-grained model for ligand-receptor interaction. *PLoS Comput Biol* 2013; 9:e1003274; PMID:24244115; <http://dx.doi.org/10.1371/journal.pcbi.1003274>